APPENDIX-I Puducherry Key Indicators

District Level Household and Facility Survey

Indicators	DLHS-3(2007-08)			DLHS-2(2002-04)				
Sample size	Total F	Rural	Urban	Total	Rural U	rban		
Households	4.000	050	0.074	4 400	0.40	0.400		
	4,333	959	3,374	4,433	943	3,488		
Ever married women (age 15-49 years)	3,851	775	3,057	NA	NA	NA		
Currently married women (age 15-44 years)	3,141	626	2,515	3,647	738	2,909		
Unmarried women (age 15-24years)	1,136	268	868	NA	NA	NA		
I. Marriage **		I		I				
Mean age at marriage for boys (marriages that occurred during the reference period)**	28.0	27.2	28.4	27.6	26.9	27.8		
Mean age at marriage for girls(marriages that occurred during the reference period)**	22.5	21.8	22.8	22.4	22.2	22.5		
Boys married below age 21 (marriages that occurred during the reference period)** (%)	3.5	5.2	2.7	2.4	6.0	1.3		
Girls married below age 18 (marriages that occurred during the reference period)** (%)	3.6	4.7	3.3	4.9	6.0	4.6		
Currently married women age 20-24 who were married before age 18 (%)	17.4	16.1	17.7	NA	NA	NA		
II. Antenatal care (based on women whose last pregnancy outcome was live/still birth during the reference period)**								
Mothers who received any antenatal check-up (%)	92.8	99.5	90.8	100.0	100.0	100.0		
Mothers who had antenatal check-up in first trimester (%)	74.8	71.0	75.9	75.5	63.8	78.7		
Mothers who had three or more ANC (%)	91.5	97.1	89.9	97.8	99.1	97.4		
Mothers who had at least one tetanus toxoid injection (%)	89.0	95.7	87.1	97.4	98.2	97.2		

APPENDIX-I Puducherry Key Indicators

District Level Household and Facility Survey

I malia a ta ma	DLHS-3(2007-08)			DLHS-2(2002-04)				
Indicators	Total Rural Urban		Total* Rural Urban					
Mothers whose Blood Pressure (BP) taken (%)	88.9	96.6	86.8	98.0	95.9	98.5		
Mothers who consumed 100 IFA Tablets (%)	56.7	45.8	60.1	31.4	28.5	32.2		
Mothers who had full antenatal check-up# (%)	47.2	40.4	49.3	30.0	28.5	30.4		
III. Delivery care (based on women whose last pregnancy outcome was live/still birth during reference period)**								
Institutional delivery (%)	99.1	97.4	99.6	97.2	98.2	96.9		
Delivery at home (%)	0.8	2.6	0.3	2.3	1.6	2.5		
Delivery at home conducted by skilled health personnel(%)	28.4	20.5	49.5	56.8	NA	NA		
Safe Delivery [®] (%)	99.4	98.0	99.9	98.5	98.2	96.9		
Mothers who received post- natal care within two weeks of delivery (%)	92.8	94.8	92.2	NA	NA	NA		
Mothers who received financial assistance for delivery under JSY ^{@@} (%)	9.8	19.5	6.8	NA	NA	NA		
IV. Child Immunization								
Number of children age 12-23 months	290	73	217	322	63	259		
Children 12-23 months fully immunized (%)	80.4	93.5	76.0	89.3	89.2	89.3		
Children 12-23 months not received any vaccination (%)	1.4	0.0	1.8	0.1	0.5	0.0		
Children 12-23 months who have received BCG vaccine (%)	96.6	100.0	95.5	98.6	99.5	98.4		
Children 12-23 months who have received 3 doses of DPT vaccine (%)	88.3	97.4	85.3	93.3	95.3	92.9		
Children 12-23 months who have received 3 doses of polio vaccine (%)	88.3	98.7	84.8	94.8	97.1	94.3		
Children 12-23 months who have received measles vaccine (%)	91.1	97.4	89.0	96.4	95.4	96.7		

APPENDIX-I Puducherry Key Indicators

District Level Household and Facility Survey

Indicators	DLHS-3(2007-08)			DLHS-2(2002-04)				
	Total F	Rural	Urban	Total*	Rural U	rban		
Children (age 9 months and above) received at least one dose of vitamin A supplement) (%)	75.9	66.8	79.2	41.9	46.6	40.5		
V. Child feeding practices (based on last-born children) (%)								
Children under 3 years breastfed within one hour of birth ⁺	70.5	67.2	71.6	73.8	77.0	72.9		
Children age 0-5 months exclusively breastfed ⁺⁺	60.6	66.0	59.0	NA	NA	NA		
Children age 6-35 months exclusively breastfed for at least 6 months	26.3	24.1	27.2	5.7	NA	NA		
Children age 6-9 months receiving solid/semi-solid food and breast milk	69.4	75.3	67.7	NA	NA	NA		
VI. Treatment of childhood diseases (based on last two surviving children born during the reference period)**								
Children with diarrhoea in the last 2 weeks who received ORS## (%)	53.8	31.7	58.3	47.7	NA	49.4		
Children with diarrhoea in the 2 weeks who were given treatment (%)	58.0	70.7	55.3	76.4	NA	73.5		
Children with acute respiratory infection or fever in last 2 weeks who were given advise or treatment (%)	84.5	81.9	85.3	90.3	NA	86.6		

Source: Ministry of Health & Family Welfare, G.O.I

- * Percentage of total (rural-urban combined) in DLHS-2 is adjusted according to rural-urban distribution of population as Pe 2001 Census.
- ** DLHS-3 reference period is form 1-1-2004 to survey date; DLHS-2 reference period is form 1-1-1999/1-1-2001 to survey date.
- # Full ANC: At least three visits for antenatal check-up, one TT injection Received and 100 IFA tablets or adequate amount of syrup consumed.

- @ Either institutional delivery or home delivery attended by skilled health Personnel (Doctor/ANM/Nurse/midwife/LHV/Other health personnel).
- @@ Janani Surasksha Yojana for deliveries form 1-4-2006 to survey date.
- + DLHS-2 data indicates children breastfed within two hours of birth.
- ++ Children who were given nothing but breast milk till the survey date.
- ## Oral Rehydration Solutions/Salts